



ISTITUTO NAZIONALE DI RICERCA METROLOGICA Repository Istituzionale

Curcumin induces apoptosis in JAK2-mutated cells by the inhibition of JAK2/STAT and mTORC1 pathways

Original

Curcumin induces apoptosis in JAK2-mutated cells by the inhibition of JAK2/STAT and mTORC1 pathways / Petiti, Jessica; Rosso, Valentina; Lo Iacono, Marco; Panuzzo, Cristina; Calabrese, Chiara; Signorino, Elisabetta; Pironi, Lucrezia; Cartellà, Antonio; Bracco, Enrico; Pergolizzi, Barbara; Beltramo, Tiziana; Fava, Carmen; Cilloni, Daniela. - In: JOURNAL OF CELLULAR AND MOLECULAR MEDICINE. - ISSN 1582-1838. - 23:6(2019), pp. 4349-4357. [10.1111/jcmm.14326]

Availability:

This version is available at: 11696/74665 since: 2022-09-22T08:35:06Z

Publisher:

WILEY

Published

DOI:10.1111/jcmm.14326

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Curcumin induces apoptosis in JAK2-mutated cells by the inhibition of JAK2/STAT and mTORC1 pathways

Jessica Petiti¹ | Valentina Rosso¹ | Marco Lo Iacono¹ | Cristina Panuzzo¹ | Chiara Calabrese¹ | Elisabetta Signorino¹ | Lucrezia Pironi¹ | Antonio Cartella¹ | Enrico Bracco² | Barbara Pergolizzi¹ | Tiziana Beltramo³ | Carmen Fava¹ | Daniela Cilloni¹

Daniela Cilloni¹

¹Department of Clinical and Biological Sciences, University of Turin, Turin, Italy

²Department of Oncology, University of Turin, Turin, Italy
³SSD Transfusional Center, San Luigi Gonzaga Hospital, Turin, Italy

Correspondence: Jessica Petiti, Department of Clinical and Biological Sciences, University of Turin, Via Carlo Cavallotti, 1, 10126 Turin, Italy.
Email: jessica.petiti@unito.it

Funding information: F.I.R.C. Fellowship for the Ricerca sul Cancro, Grant/Award Number: 10005

Myeloproliferative neoplasms are chronic myeloid cancers divided in Philadelphia positive and negative. The JAK2 V617F is the most common mutation in Philadelphia negative patients and results in a constitutive activation of the JAK/STAT pathway, conferring a proliferative advantage and apoptosis inhibition. Recent studies identified a functional crosstalk between the JAK/STAT and mTOR pathways. The identification of an effective therapy is often difficult, so the availability of new therapeutic approaches might be attractive. Previous studies showed that curcumin, the active principle of the *Curcuma longa*, can suppress JAK2/STAT pathways in different type of cancer and injuries. In this study, we investigated the anti-proliferative and pro-apoptotic effects of curcumin in JAK2 V617F-mutated cells. HEL cell line and cells from patients JAK2 V617F mutated have been incubated with increasing concentrations of curcumin for different time. Apoptosis and proliferation were evaluated. Subsequently, JAK2/STAT and AKT/mTOR pathways were investigated at both RNA and protein levels. We found that curcumin induces apoptosis and inhibition of proliferation in HEL cells. Furthermore, we showed that curcumin inhibits JAK2/STAT and mTORC1 pathways in JAK2 V617F-mutated cells. This inhibition suggests that curcumin could represent an alternative strategy to be explored for the treatment of patients with myeloproliferative neoplasms.

KEYWORDS
curcumin, JAK/STAT, JAK2 V617F, mTORC1, Myeloproliferative neoplasms

7 HUPV RI XVH

7 KLV DUWLFQH LV PDGHV FULSWLRQ LQ WKH

1 | INTRODUCTION

Myeloproliferative neoplasms (MPNs) are clonal hematopoietic stem cell disorders characterized by an uncontrolled proliferation of one or more elements of the myeloid lineage.¹ In addition to chronic

myeloid leucemia (CML), characterized by the presence of the Philadelphia chromosome, the Philadelphia negative MPNs are essential thrombocythemia (ET), polycythemia vera (PV) and primary myelofibrosis (PMF). The JAK2 V617F is the most frequent mutation in patients affected by PV, ET and PMF. JAK2 gene encodes for a

Jessica Petiti and Valentina Rosso contributed equally to this manuscript.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 John Wiley & Sons Ltd and Foundation for Cellular and Molecular Medicine. *J Cell Mol Med.* 2019;23:4349–4357.

non-receptor tyrosine kinase crucial for signal transduction downstream of the erythropoietin, thrombopoietin and related receptors that control erythrocyte and megakaryocyte expansion.² Activated JAK2 phosphorylates STATs proteins, specifically STAT5 and STAT3, that homodimerize and translocate to the nucleus. Activated STATs induce the expression of target genes, such as PIM-1, PIM-2 and PIM-3, serine/threonine kinases that promote cells survival, proliferation and therapy resistance.³ JAK2 mutations directly activate JAK/STAT signalling and make myeloproliferation cytokine independent or hypersensitive. JAK/STAT deregulation is critical for MPNs developing and progression. Furthermore, recent studies identified the role of mTOR pathway in MPNs, highlighting a functional crosstalk between the JAK/STAT and mTOR.^{4,5} mTOR is a serine/threonine kinase that regulates cellular metabolism, growth and survival and it may form different proteins complexes: mTORC1 and mTORC2. mTORC1 is composed of mTOR, Raptor, GβL and DEPTOR and it is regulated by AKT. In normal cells, mTORC1 is essential for erythroid and megakaryocytic differentiation through the activation of downstream effectors including 4eBP1 and p70s6K.⁶ This pathway has been found deregulated particularly in megakaryocytes of MPNs patients.⁷ The deregulation of JAK/STAT and mTOR pathways induces an inflammatory state with aberrant cytokine expression.⁸ Given the heterogeneous clinical needs of MPNs patients, determination of a standard therapeutic protocol is often difficult. Moreover, targeted therapy with JAK inhibitors revealed to have some limits in terms of efficacy,⁹ therefore it is necessary to find additional approaches to improve the results so far obtained.

Curcumin is the active phytochemical component isolated from the rhizome of the *Curcuma longa* plant. Curcumin is a highly pleiotropic molecule with multiple pharmacological effects, such as anti-inflammatory, anti-microbial, anti-oxidative and anti-proliferative activities.^{10,11} Extensive preclinical trials have indicated curcumin therapeutic potential against a wide range of human diseases.¹² Previous studies showed that curcumin can suppress JAK2/STAT signalling pathways in different type of cancer and injuries.^{13,14} Chen et al demonstrated that curcumin increased the transcript levels of SOCS-3, an important negative regulator of JAK2, and significantly inhibited the clonogenic activity of hematopoietic progenitors from MPNs patients.¹⁵ Furthermore, curcumin was able to dissociate Raptor from mTOR, by inhibiting mTORC1 signalling and the phosphorylation of its downstream effectors in different cell lines.¹⁶ In this study, we investigated the effect of curcumin on JAK2 V617F cell line and in primary cells from MPNs patients. Our results suggest that curcumin inhibits proliferation and activates cell death program by modulating JAK2/STAT and mTORC1 pathways.

2 | MATERIALS AND METHODS

2.1 | Cells culture conditions

HEL cell line was purchased from American Type Culture Collection (ATCC, Manassas, USA). HEL cells were grown in RPMI 1640 medium supplemented with 200 nmol/L Glutamine (EuroClone), 10%

inactivated foetal bovine serum (FBS, Sigma-Aldrich) and 0.1% penicillin/streptomycin and maintained at 37°C with 5% CO₂.

2.2 | Patients cohort

After informed consent, human peripheral blood (PB) leucocytes were isolated by Buffy Coat procedure from 30 MPNs patients (24 were PV, 4 ET and 2 PMF; the median age was 63 years (range 20-86); 18 were males and 12 females) and 10 healthy donors. All samples obtained from patients were JAK2 V617F mutated. The study was approved by the ethic committee on 16 December 2015 (number of approval 212/2015).

2.3 | Cells treatment

HEL cells were incubated with different concentrations (10, 15, 20, 30 μmol/L) of curcumin (stock solution 50 mmol/L in DMSO, #C1386, Sigma-Aldrich) for 24 and 48 hours. Leucocytes isolated from MPNs patients were incubated with 30 μmol/L of curcumin in IMDM (EuroClone) supplemented with 20% inactivated FBS for 20 hours. After incubation, apoptosis and proliferation were evaluated and total RNA and proteins were extracted as described below.

2.4 | Apoptosis and viability assays

Apoptosis was evaluated using APC Annexin V (BioLegend), according to the manufacturer's instructions. Cells were analysed by flow cytometry (FACS) and the apoptotic fraction was defined as annexin V positive. HEL cells were treated with increasing concentration of curcumin for 24 and 48 hours. After incubation, cells were harvested, resuspended in Flow Cytometry Staining Buffer (#FC001, R&D Systems) and labelled with 1 μg/μL propidium iodide (PI). Live cells (PI negative fraction) were counted by FACS.

2.5 | Protein extraction and immunoblotting

To perform immunoblotting analysis, we used human HEL cells and cells from MPNs patients. To isolate total protein content, samples were lysed in ice with Ripa buffer (50 mmol/L Tris-HCl pH 8.0, 150 mmol/L NaCl, 1% NP40, 0.5% DOC, 0.1% SDS with freshly added of protease and phosphatase inhibitors cocktail [Sigma-Aldrich]) and cell debris were removed by centrifugation at 14 000g at 4°C for 15 minutes. Protein concentration was determined by Bio-Rad Protein Assay (Bio-Rad): 30 μg of each total cell lysate were loaded, resolved through SDS-PAGE 6%-12% gel and electroblotted onto 0.2 μm nitrocellulose membranes (Bio-Rad). After blocking with 5% BSA (Sigma-Aldrich) in TBS 1x plus 0.2% Tween-20 (Sigma-Aldrich) for at least 1 hour at RT, membranes were incubated overnight (ON) at 4°C with the primary antibodies listed in Table 1 with a dilution of 1:1000. As a secondary antibody, we used goat antimouse IgG-HRP (sc-2005), mouse anti-goat IgG-HRP (sc-2354) and goat anti-rabbit IgG-HRP (sc2004) [Santa Cruz Biotechnology] in a dilution of 1:7000 for 1 hour at RT. Immunoreactive bands were visualized by Clarity Western ECL Substrate (Bio-Rad). Quantification was performed using Image Lab program (Bio-Rad).

concentrations. In addition, the curcumin growth inhibitory effect was dose and time-dependent, reaching the maximum effect at the dose of 20 $\mu\text{mol/L}$ for 48 hours, by reducing the proliferation of 93% (Figure 1A). The results obtained evaluating apoptosis with Annexin V integrate and confirm viability analysis. In particular, we showed that the percentage of apoptotic cells increase of 2.1, 3.6, 5.0 and 5.9 times at 24 hours and 2.1, 4.7, 6.3 and 8.6 at 48 hours, respectively after treatment with 10, 15, 20, and 30 $\mu\text{mol/L}$ of curcumin (Figure 1B). Furthermore, to investigate the intrinsic apoptosis at the protein level, treated HEL cells were harvested at 24 hours and analysed by Western blot. As shown in Figure 1C, the protein expression of cleaved caspase-3 was markedly increased in the curcumin-treated HEL cells according to drug concentration. In particular, a curcumin concentration of 30 $\mu\text{mol/L}$ showed a sharp increase of cleaved caspase-3. This data confirmed the FACS analysis and indicated that curcumin induced apoptosis in a dose-dependent manner in JAK2 V617F-mutated cell line.


3.2 | Curcumin affects JAK2/STAT pathway in HEL cell line

To investigate the modulation of JAK/STAT pathway, HEL cells were treated with various concentrations of curcumin (0-30 $\mu\text{mol/L}$) for 24 hrs and JAK2, STAT3 and STAT5 phosphorylation status was evaluated by immunoblotting. Data showed that curcumin reduced the tyrosine phosphorylation of JAK2, STAT3 and STAT5 in a dose-dependent manner. Although the cells treated with curcumin 20 and 30 $\mu\text{mol/L}$ showed a reduction of JAK2 phosphorylation of about 30%, the functionality of JAK2 to phosphorylate its effectors was

strongly compromised. The phosphorylation of STAT5 and STAT3, the direct effectors of JAK2, was strongly inhibited by curcumin. Indeed, we observed that phosphorylation of STAT5 was 60% reduced, while STAT3 was extremely sensitive to curcumin-mediated JAK2 inhibition and its phosphorylation decreased to 20% even at the lower curcumin dose and was completely blocked at higher concentrations. Furthermore, the protein expression of SOCS-1 and SOCS-3, negative regulators of JAK/STAT pathway, resulted clearly up-regulated after curcumin treatment. Indeed, their expression increased of three and 1.4 times, respectively, even at the lower curcumin dose and of seven and two times, respectively, at higher concentrations (Figure 2A). A similar behaviour was observed also at mRNA level; indeed, curcumin significantly increased the expression of SOCS-1 and SOCS-3, displaying an up-regulation of about six times even at 15 $\mu\text{mol/L}$ of curcumin ($P < 0,01$) (Figure 2B). These results were further corroborated by measuring the transcriptional activity of JAK downstream effectors STATs on mRNA expression of PIM family members. Curcumin significantly decreased the expression of PIM1, PIM2 and PIM3, showing a down-modulation of about two times compared to control when cells were treated with a concentration higher than 20 $\mu\text{mol/L}$ (PIM1 and 3 $P < 0.01$ and PIM2 $P < 0.05$) (Figure 2B).

3.3 | Curcumin modulates mTORC1 signalling by the inhibition of PDK and AKT in HEL cells line

To investigate how curcumin affects mTORC1 pathway, HEL cells were treated with various concentrations of curcumin (0-30 $\mu\text{mol/L}$)



& ! • JHEL cells treated with different concentration of curcumin (0-30 $\mu\text{mol/L}$) for 24 and 48 h. A, Effect of curcumin on cell viability. The percent of live cells was calculated by normalizing with the control ($n = 5$). B, Effect of curcumin on apoptosis ($n = 5$). C, Caspase-3 Cleaved expression was evaluated by SDS-PAGE. The GAPDH level was used as loading control. The relative intensity of each band is shown under the blot as fold change (FC) compared to untreated control, to which a value of 1 unit was assigned. Statistical analyses were performed using the two-tailed Mann-Whitney U test, comparing conditions two by two respect to not treated condition. All the analysis with confidence level major of 95% are indicated like significant and marked as followed: * $P \leq 0.05$; ** $P \leq 0.01$.


for 24 hours and analysed by immunoblotting. Our results indicated that curcumin affects the principal modulator of mTORC1: AKT. Phosphorylation of AKT in Thr308 was reduced in dose-dependent manner and its principal activator PDK was inhibited by low dose curcumin and it appeared unphosphorylated at the maximum concentration utilized (Figure 3). The consequence was the reduction of the complex between mTOR and Raptor, as observed in Figure 3, especially at the highest drug concentration. Furthermore, by analysing the signal cascade, we observed the deregulation of the principal downstream effectors of mTORC1. In particular, we showed the dephosphorylation of p70s6K, which is usually activated by


phosphorylation, and the increase of unphosphorylated form of 4eBP1, usually inhibited through mTORC1 phosphorylation.

3.4 | Curcumin induces apoptosis and reduces PIM family members and CD177 expression in JAK2 mutated MPNs patients

Due to the promising results in vitro, we tried to evaluate the effect of curcumin also in patient specimens. For these purposes, leucocytes from JAK2 mutated patients were treated with curcumin at the highest concentration utilized in vitro (30 µmol/L) for 20 hours and the

JHEL cells treated with different concentration of curcumin (0-30 µmol/L) for 24 h. A, The indicated proteins of JAK2/STAT5 signalling were detected by SDS-PAGE. Representative Western blots were shown (n = 3). The relative intensity of each band is shown under each blot as fold change (FC) compared to untreated control. B, The mRNA expression levels of PIM family members, Socs-1 and Socs-3 were evaluated by qRT-PCR. Results are expressed as $-\Delta\Delta Ct$ (n = 7). Statistical analyses were performed using the two-tailed Mann-Whitney U test, comparing conditions two by two respect to not treated condition. All the analysis with confidence level major of 95% are indicated like significant and marked as followed: *P ≤ 0.05; **P ≤ 0.01.





& ! ' JHEL cells treated with different concentrations of curcumin (0-30 μmol/L) for 24 h. Phosphorylation status of PDK1 and AKT (n = 4), co-immunoprecipitation of mTOR-Raptor complex (n = 3) and phosphorylation status of p70s6k and 4EBP1 (n = 4) were evaluated by SDS-PAGE. The relative intensity of each band is shown under each blot as the fold change (FC) compared to untreated control. The analysed pathways were indicated on right side


expression of PIM family members and CD177 were evaluated by qRT-PCR. The results were in accordance with in vitro analysis and showed that PIM1, PIM2, PIM3 and CD177 mRNA expression was significantly down-regulated in cells treated with curcumin compared to not treated samples (PIM1 $P = 0.0014$; PIM2 $P = 0.0073$; PIM3 $P = 0.011$; CD177 $P < 0.0001$) (Figure 4A). Interestingly, except for PIM2 ($P = 0.025$) (Figure 4A), the mRNA expression of PIM family members and CD177 after treated return to the levels of healthy subjects (PIM1 $P = 0.1$; PIM3 $P = 0.07$ and CD177 $P = 0.58$). Furthermore, the expression of cleaved caspase-3, evaluated by SDS-PAGE, was significantly up-regulated in patients' cells treated with curcumin ($P < 0.001$), while it has not changed in cells from healthy donors (Figure 4B). These results suggest that curcumin treatment affects JAK/STAT pathway and induces apoptosis in cells from JAK2 V617F mutated patients.

4 | DISCUSSION

The majority of Philadelphia negative MPNs shows the single base JAK2 V617F mutation that results in a constitutive activation of the

JAK/STAT pathway, conferring a proliferative advantage and apoptosis inhibitions. The discovery of JAK2 mutations allowed the development of JAK2 inhibitors for target therapies, including ruxolitinib, approved for patients with primary or secondary myelofibrosis and hydroxyurea-resistant or intolerant PV patients. Although ruxolitinib has brought enormous benefits to patients, evidence that JAK2 inhibition can cure the disease, at least in PMF, remains debated.^{18,19} The difficulty in treating patients affected by MPNs encourages the identification of new effective drugs, including some natural compounds, like curcumin. Curcumin has been found to exert its anti-inflammatory and anti-carcinogenic effects by targeting JAK2 signalling in different type of neoplasms and injuries,^{14,20} but the effects of this phytochemical on JAK2-mutated cells have been poorly studied until now.

In this work, we investigated, at both RNA and protein levels, the effects of curcumin on JAK2/STAT pathway, known to be altered in MPNs patients. Our data are in accordance to what has been already published by Chen and colleagues, who demonstrated that curcumin up-regulate SOCS-1 and SOCS-3 which are important negative regulators of JAK2.¹⁵ In addition, we showed that curcumin strongly inhibited the proliferation and induced apoptosis in



& ! “ JLeucocytes of MPNs patients (pts) were treated with 30 μmol/L curcumin for 20 h. A, The mRNA expression of PIM family members and CD177 was evaluated by qRT-PCR in MPNs pts and healthy controls. Results are expressed as $-\Delta Ct$ ($n = 20$). Statistical analyses were performed using paired t test to compare treated and not treated MPNs patients' cells and the two-tailed Mann-Whitney U test to compare treated patients with healthy donors' cells. All the analysis with confidence level major of 95% are indicated like significant and marked as followed: $*P \leq 0.05$; $**P \leq 0.01$; $***P \leq 0.001$. B, Caspase-3 Cleaved expression was evaluated by SDS-PAGE. The GAPDH level was used to normalize data. Representative Western blots of three patients and three healthy controls were shown

a dose and time-dependent manner in HEL cells. Furthermore, we found that curcumin markedly reduced JAK2 phosphorylation and, consequently, the activation of its downstream effectors STAT3 and STAT5 in HEL cells, according to the results obtained by Zhu et al in severe acute pancreatitis¹⁴ and by Kim et al in rat primary microglia.²⁰ The inhibition of JAK2/STAT pathway, observed in our study, induced the down-regulation of PIM family members, PIM1, PIM2 and PIM3 in HEL cell line and in patient specimens. These small kinases are known to be involved in leukemogenesis and in ruxolitinib resistance in MPNs cells.^{21,22} As Hitosugi et al identified that constitutively activated JAK2 regulates AKT through the phosphorylation of PDK1,²³ we focused our attention on this pathway. Our data showed that curcumin-mediated inhibition of JAK2 reduces the activation of PDK1 and, consequently, the Thr-308 AKT phosphorylation in HEL cells. It has been reported that AKT directly regulates mTORC1 signalling,²⁴ so we investigated the effect of curcumin on this pathway. In agreement with the results obtained by Beevers et al in human rhabdomyosarcoma cells, our results in HEL cells showed that curcumin negatively regulated the mTORC1 complex formation. As a consequence, we observed a strong inhibition of mTORC1 downstream effectors 4eBP1 and p70s6K phosphorylation, both involved in cell growth and proliferation.¹⁶ This data are in accordance with the hypothesis that mTORC1 is a JAK2 downstream pathway and it is involved in MPNs pathogenesis.²⁵ Due to the promising data

obtained in cell line, we tried to evaluate the effect of curcumin also in patient samples. Our preliminary data showed that the in vitro curcumin treatment of cells from MPNs patients significantly reduced the expression of PIM family members, bringing back their mRNA expressions to comparable levels with healthy subjects ones. The curcumin effect was strengthened by the observation that CD177, an antigen over expressed in neutrophils of the majority of MPNs patients, was down-regulated by curcumin.²⁶

Ishida et al demonstrated that HEL cells are only partially dependent on JAK2 V617F for survival and this may explain the very limited effect of ruxolitinib and its inefficiency to eradicate JAK2-mutated clone in MPNs patients.²⁷ Interestingly, in this study, we observed that curcumin affects both proliferation and survival of HEL cells, suggesting that its role in MPNs could be more effective in blocking neoplastic cells respect to the common JAK2 inhibitors, such as ruxolitinib. In vitro studies support this hypothesis by indicating that co-treatment of mTOR and JAK2 inhibitors resulted in synergistic activity against JAK2 V617F-mutated cells,⁵ although mTORC1 inhibitors can cause numerous side effects that could prevent their use.²⁸ In contrast, curcumin turned out to be safe and non-toxic in many different trials,^{29,30} so its use for MPNs treatment could represent an excellent alternative to common mTORC1 inhibitors. Although the curcumin employment for MPNs treatment needs to be further investigated, our preclinical data encourage its use for the future therapies.

In conclusion, this study showed that curcumin exerts an antitumor effect on human JAK2-mutated cells by inducing apoptosis and inhibition of proliferation, through the regulation of both JAK2/STAT and mTORC1 pathways. These findings suggest that curcumin seems to be a promising nutraceutical compound that should be further evaluated in different pharmaceutical formulation for the treatment of MPNs.

ACKNOWLEDGEMENTS

The study was funded by grants from AIRC. The study was partially funded by a special grant from 'AIRC 5 per mille' to the AGIMM group (AIRC-Gruppo Italiano Malattie Molecolari); for a complete list of AGIMM investigators see at <http://www.progettoagimm.it>.

CONFLICT OF INTEREST

The authors declare that they have no competing interests.

AUTHOR CONTRIBUTIONS

JP and VR designed the study, performed the experiments and wrote the manuscript. ML analysed data and revised the manuscript. CP performed FACS analysis. CC, ES, LP and AC collected samples. EB and BP supervised the experiments. TB provided samples. CF and DC supervised the experiments and wrote the manuscript. All authors read and approved the final manuscript.

ORCID

Jessica Petiti  <https://orcid.org/0000-0001-8640-2462>

Daniela Cilloni  <https://orcid.org/0000-0001-6346-4791>

REFERENCES

- Mithraprabhu S, Grigoriadis G, Khong T, Spencer A. Deacetylase inhibition in myeloproliferative neoplasms. *Invest New Drugs*. 2010;28(Suppl 1):S50-S57.
- Levine RL, Gilliland DG. Myeloproliferative disorders. *Blood*. 2008;112:2190-2198.
- Jatiani SS, Baker SJ, Silverman LR, Reddy EP. Jak/STAT pathways in cytokine signaling and myeloproliferative disorders: approaches for targeted therapies. *Genes & cancer*. 2010;1:979-993.
- Bartalucci N, Calabresi L, Balliu M, et al. Inhibitors of the PI3K/mTOR pathway prevent STAT5 phosphorylation in JAK2V617F mutated cells through PP2A/CIP2A axis. *Oncotarget*. 2017;8:96710-96724.
- Bogani C, Bartalucci N, Martinelli S, et al. mTOR inhibitors alone and in combination with JAK2 inhibitors effectively inhibit cells of myeloproliferative neoplasms. *PLoS ONE*. 2013;8:e54826.
- Martelli AM, Evangelisti C, Chiarini F, et al. The emerging role of the phosphatidylinositol 3-kinase/Akt/mammalian target of rapamycin signaling network in normal myelopoiesis and leukemogenesis. *Biochem Biophys Acta*. 2010;1803:991-1002.
- Vicari L, Martinetti D, Buccheri S, et al. Increased phospho-mTOR expression in megakaryocytic cells derived from CD34+ progenitors of essential thrombocythaemia and myelofibrosis patients. *Br J Haematol*. 2012;159:237-240.
- Quintas-Cardama A, Verstovsek S. Molecular pathways: Jak/STAT pathway: mutations, inhibitors, and resistance. *Clin Cancer Res*. 2013;19:1933-1940.
- Tefferi A. JAK inhibitors for myeloproliferative neoplasms: clarifying facts from myths. *Blood*. 2012;119:2721-2730.
- Karunakaran D, Rashmi R, Kumar TR. Induction of apoptosis by curcumin and its implications for cancer therapy. *Curr Cancer Drug Targets*. 2005;5:117-129.
- Sharma RA, Gescher AJ, Steward WP. Curcumin: the story so far. *Eur J Cancer*. 2005;41:1955-1968.
- Aggarwal BB, Kumar A, Bharti AC. Anticancer potential of curcumin: preclinical and clinical studies. *Anticancer Res*. 2003;23:363-398.
- Wu L, Guo L, Liang Y, Liu Xu, Jiang L, Wang L. Curcumin suppresses stem-like traits of lung cancer cells via inhibiting the JAK2/STAT3 signaling pathway. *Oncol Rep*. 2015;34:3311-3317.
- Zhu S, Zhang C, Weng Q, Ye B. Curcumin protects against acute renal injury by suppressing JAK2/STAT3 pathway in severe acute pancreatitis in rats. *Exp Ther Med*. 2017;14:1669-1674.
- Chen CQ, Yu K, Yan QX, et al. Pure curcumin increases the expression of SOCS1 and SOCS3 in myeloproliferative neoplasms through suppressing class I histone deacetylases. *Carcinogenesis*. 2013;34:1442-1449.
- Beevers Cs, Chen L, Liu L, Luo Y, Webster Nj, Huang S. Curcumin disrupts the Mammalian target of rapamycin-raptor complex. *Can Res*. 2009;69:1000-1008.
- Peterson TR, Laplante M, Thoreen CC, et al. DEPTOR is an mTOR inhibitor frequently overexpressed in multiple myeloma cells and required for their survival. *Cell*. 2009;137:873-886.
- Barosi G, Zhang MJ, Gale RP. Does ruxolitinib improve survival of persons with MPN-associated myelofibrosis? should it? *Leukemia*. 2014;28:2267-2270.
- Cervantes F, Pereira A. Does ruxolitinib prolong the survival of patients with myelofibrosis? *Blood*. 2017;129:832-837.
- Kim HY, Park EJ, Joe EH, Jou I. Curcumin suppresses Janus kinase-STAT inflammatory signaling through activation of Src homology 2 domain-containing tyrosine phosphatase 2 in brain microglia. *J Immunol*. 2003;171:6072-6079.
- Saurabh K, Scherzer MT, Shah PP, et al. The PIM family of oncoproteins: small kinases with huge implications in myeloid leukemogenesis and as therapeutic targets. *Oncotarget*. 2014;5:8503-8514.
- Mazzacurati L, Lambert QT, Pradhan A, Griner LN, Huszar D, Reuther GW. The PIM inhibitor AZD1208 synergizes with ruxolitinib to induce apoptosis of ruxolitinib sensitive and resistant JAK2-V617F-driven cells and inhibit colony formation of primary MPN cells. *Oncotarget*. 2015;6:40141-40157.
- Hitosugi T, Fan J, Chung T-W, et al. Tyrosine phosphorylation of mitochondrial pyruvate dehydrogenase kinase 1 is important for cancer metabolism. *Mol Cell*. 2011;44:864-877.
- Hahn-Windgassen A, Nogueira V, Chen C-C, Skeen JE, Sonenberg N, Hay N. Akt activates the mammalian target of rapamycin by regulating cellular ATP level and AMPK activity. *J Biol Chem*. 2005;280:32081-32089.
- Skoda RC, Duek A, Grisouard J. Pathogenesis of myeloproliferative neoplasms. *Exp Hematol*. 2015;43:599-608.
- Stroncek DF, Caruccio L, Bettinotti M. CD177: A member of the Ly-6 gene superfamily involved with neutrophil proliferation and polycythemia vera. *J Transl Med*. 2004;2:8.
- Ishida S, Akiyama H, Umezawa Y, et al. Mechanisms for mTORC1 activation and synergistic induction of apoptosis by ruxolitinib and BH3 mimetics or autophagy inhibitors in JAK2-V617F-expressing leukemic cells including newly established PVTL-2. *Oncotarget*. 2018;9:26834-26851.

28. Pallet N, Legendre C. Adverse events associated with mTOR inhibitors. *Expert Opin Drug Saf.* 2013;12:177-186.
29. Aggarwal ML, Chacko KM, Kuruvilla BT. Systematic and comprehensive investigation of the toxicity of curcuminoid essential oil complex: a bioavailable turmeric formulation. *Mol Med Rep.* 2016;13:592-604.
30. Chainani-Wu N. Safety and anti-inflammatory activity of curcumin: a component of tumeric (*Curcuma longa*). *J Altern Complement Med.* 2003;9:161-168.

How to cite this article: Petiti J, Rosso V, Lo Iacono M, et al. Curcumin induces apoptosis in JAK2-mutated cells by the inhibition of JAK2/STAT and mTORC1 pathways. *J Cell Mol Med.* 2019;23:4349-4357. <https://doi.org/10.1111/jcmm.14326>