



ISTITUTO NAZIONALE DI RICERCA METROLOGICA Repository Istituzionale

The time as an emergent property of quantum mechanics, a synthetic description of a first experimental approach

Original

The time as an emergent property of quantum mechanics, a synthetic description of a first experimental approach / Moreva, Ekaterina; Brida, Giorgio; Gramegna, Marco; Giovannetti, V.; Maccone, L.; Genovese, Marco. - In: JOURNAL OF PHYSICS. CONFERENCE SERIES. - ISSN 1742-6588. - 626:(2015), pp. 012019.012019-1-012019.012019-7. [0.1088/1742-6596/626/1/012019]

Availability:

This version is available at: 11696/34002 since:

Publisher:

Published

DOI:0.1088/1742-6596/626/1/012019

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

The time as an emergent property of quantum mechanics, a synthetic description of a first experimental approach

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 J. Phys.: Conf. Ser. 626 012019

(<http://iopscience.iop.org/1742-6596/626/1/012019>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

This content was downloaded by: mgramagna

IP Address: 193.204.114.88

This content was downloaded on 02/09/2015 at 16:07

Please note that [terms and conditions apply](#).

The time as an emergent property of quantum mechanics, a synthetic description of a first experimental approach

E Moreva^{1,2}, G Brida¹, M Gramegna¹, V Giovannetti³, L Maccone⁴
and M Genovese^{1*}

¹INRIM, strada delle Cacce 91, 10135 Torino, Italy

²International Laser Center of M.V.Lomonosov Moscow State University, 119991, Moscow, Russia

³NEST, Scuola Normale Superiore and Istituto Nanoscienze-CNR, piazza dei Cavalieri 7, I-56126 Pisa, Italy

⁴Dip. Fisica “A. Volta”, INFN Sez. Pavia, Univ. of Pavia, via Bassi 6, I-27100 Pavia, Italy

E-mail: *m.genovese@inrim.it

Abstract. The “problem of time” in present physics substantially consists in the fact that a straightforward quantization of the general relativistic evolution equation and constraints generates for the Universe wave function the Wheeler-De Witt equation, which describes a static Universe. Page and Wootters considered the fact that there exist states of a system composed by entangled subsystems that are stationary, but one can interpret the component subsystems as evolving: this leads them to suppose that the global state of the universe can be envisaged as one of this static entangled state, whereas the state of the subsystems can evolve. Here we synthetically present an experiment, based on PDC polarization entangled photons, that shows a practical example where this idea works, i.e. a subsystem of an entangled state works as a “clock” of another subsystem.

1. Introduction

“...an infinite series of times, in a dizzily growing, ever spreading network of diverging, converging and parallel times. This web of time—the strands of which approach one another, bifurcate, intersect or ignore each other through the centuries—embraces every possibility.” [Borges]

“Quid est ergo tempus? si nemo ex me quaerat, scio; si quaerenti explicare velim, nescio.”

[1] As Augustinus Hipponensis present physicists are in the situation where time is an essential physical parameter whose meaning is intuitively clear, but several problems arise when they try to provide a clear definition of time. First of all the definition of time is different in different branches of physics as classical and non-relativistic quantum mechanics (a fixed background parameter), special relativity (a proper time for each observer, time as fourth coordinate; set of privileged inertial frames) or general relativity (time is a general spacetime coordinate: not at all an absolute time, time non orientability, closed timelike curves)[2]. More specifically, the problem of time [2, 3, 4, 5, 6, 7, 8, 9, 10, 11] in physics consists in the fact that a straightforward quantization of the general relativistic evolution equation and constraints generates the Wheeler-



De Witt equation $H_{tot}|\Psi\rangle = 0$, where H_{tot} is the global Hamiltonian of the universe and $|\Psi\rangle$ is its state. Obviously, this means that the state of the universe must be static, which clashes with our everyday experience of an evolving world.

Apart from the Wheeler-De Witt equation, one can also consider that a time shift of the state of the whole universe must be unobservable from a purely physical consideration: if shifting the state of the whole universe, there is nothing external that can keep track of this shift, so this shift must be unobservable (it is a trivial application of Mach's principle). Same considerations would apply, of course, also to the spatial degrees of freedom: a global shift in position of the whole state of the universe must similarly be unobservable, so the universe must be in an eigenstate of its momentum operator. In the following we will only consider temporal degrees of freedom.

Incidentally, this idea of a static universe surprisingly re-proposes, of course rephrased in modern language, ideas stemming from the work of Parmenides of Elea [”the phenomena of movement and change are simply appearances of a static, eternal reality”] and then diffused in roman-hellenistic word [”Tempus item per se non est, sed rebus ab ipsis consequitur sensus, transactum quid sit in aevo, tum quae res instet, quid porro deinde sequatur”¹ Titus Lucretius Carus, De Rerum Natura].

The Page and Wootters scheme [12, 13], which developed some previous ideas [14], is based on the fact that there exist states of a system composed by entangled subsystems that are stationary, but one can interpret the component subsystems as evolving. One can then suppose that the global state of the universe is one of this static entangled state, whereas the state of the subsystems (us, for example) can evolve. This solves in an extremely elegant way the problem of time. Incidentally, Page and Wootters' proposal naturally embodies the philosophy of relationalism [23, 24] and operationalism, since time is only defined in relation to clocks and to its measurement procedure [33, 34].

Up to now, all these considerations were of theoretical character. Here we epitomize an experimental approach [37] to this problem by providing an emblematic example of Page and Wootters idea at work, visualizing how time could emerge from a static (with respect to an abstract external time) entangled state. Even though the total state of a system is static, time is recovered as correlations between a subsystem that acts as a clock and the rest of the system that evolves according to such a clock. We use a system composed of two entangled photons (a paradigmatic system for several emblematic experiments [35]): the rotation of the polarization of the first acts as a clock for proving an evolution of the polarization of the second. Nonetheless, we demonstrate that the joint polarization state of both photons does not evolve. The Page-Wootters mechanism has been criticised [4, 25], and Gambini et al. have proposed some new ideas [21, 31] to overcome these criticisms. Our experiment demonstrates also some aspects of these ideas.

2. The experiment

The experimental setup is schematically depicted in Figure 1. It consists of two blocks, ”preparation” and ”measurement”. The preparation block allows producing a family of ququarts biphoton polarization states [36] of the form:

$$|\Psi\rangle = \cos\theta |HH\rangle + e^{i\varphi} \sin\theta |VV\rangle \quad (1)$$

It includes two orthogonally oriented BBO crystals (1mm) that, pumped by a 700 mW cw Ar laser operating at 351 nm, generate a pair of the basic ququart states via type-I spontaneous parametric down conversion in collinear, frequency degenerate regime around the central wavelength of 702 nm. The basic state amplitudes are controlled with the help of a

¹ Time also exists not by itself, but simply from the things which happen the sense apprehends what has been done in time past, as well as what is present and what is to follow after




Figure 1. Scheme for the two experiments. The experimenter in observer mode (block a) can prove the time evolution of the first photon (upper path) using only correlation measurements between it and the clock photon (lower path) without access to an external clock. The experimenter in super-observer mode (block b) proves that the global state of the system is static through state tomography. The blue boxes (A) represent different thicknesses of birefringent plates which evolve the photons by rotating their polarization: different thicknesses represent different time evolutions. The dashed box (B) represents a (known) phase delay of the clock photon only; PBS stands for polarizing beam splitter in the H/V basis; BS for beam splitter.

Thompson prism (V), oriented vertically and the half-wave plate $\lambda/2$ at an angle θ . Two 1 mm quartz plates (QP), that can be rotated along the optical axis, introduce a phase shift φ between horizontally and vertically polarized type-I entangled [40] biphotons. Non-linear crystals and quartz plates are placed into a temperature-stabilized closed box for achieving stable phase-matching conditions all the time. The preparation block also includes a non-polarizing beam splitter (BS), which is used to split the initial (collinear) biphoton field into two spatial modes 1, 2. For preparing the singlet Bell state

$$|\Psi^-\rangle = \frac{1}{\sqrt{2}} (|H_1\rangle |V_2\rangle - |V_1\rangle |H_2\rangle) \quad (2)$$

parameters $\theta = 45^\circ, \varphi = 0^\circ$, were selected and an additional half-wave plate $\lambda/2$ at 45° was introduced in the transmitted arm.

The measurement part can be mounted in two different modes "observer" and "super-observer". "Observer" mode (Figure 1, block a) serves as proving that one subsystem (polarization of the upper photon) evolves with respect to a clock constituted by the other subsystem (polarization of the lower photon). In the "super-observer" mode ((Figure 1, block b)) we show that the global state of the two photons is static with respect to any "external" clock. Let us analyze the observer mode first.

"Observer" mode: Each arm of a measurement block contains interference filters (IF) with central wavelength 702 nm (FWHM 1 nm) and polarizing beam splitter (PBS). Four avalanche photodiode detectors (APD) are placed at the ends and connected to a coincidence count scheme (CC) with 1.5 ns time window.

In this mode the polarization of the photon in the lower arm is used as a clock for proving an evolution of the polarization of the photon in the upper arm. In other words, the first polarizing beam splitter PBS acts as a non-demolition measurement that measures (in the H/V basis)

the polarization of the second photon in the state $|\Psi^-\rangle$ (2). Once it is initialized in the H state, the polarization of both photons evolves in the birefringent quartz plates (A) at 45° as $|H\rangle \rightarrow \frac{1}{\sqrt{2}}(|H\rangle + e^{i\delta}|V\rangle)$, $|V\rangle \rightarrow \frac{1}{\sqrt{2}}(|H\rangle - e^{i\delta}|V\rangle)$, where δ is the materials optical thickness. The clock readout is performed on the lower detectors, which measure whether the photon is $|H\rangle$ or $|V\rangle$. This clock is extremely primitive, since it has a dial with only two values: either it found with H polarization, corresponding to time t_1 or with V polarization, corresponding to time t_2 .

In the "observer" mode we do not have access to any external clock, the only information we obtain are correlations (coincidences) between the detectors. Thus, this primitive clock allows extracting the information about the polarization of the upper photon in a defined state, for example $|V\rangle$, only from registering coincidence counts between the detectors 1 – 3 (that corresponds to the measurement at the moment time t_1) or 2-3 (that corresponds to time t_2). Furthermore, one does not have any knowledge about the past time after the clock initialization (knowledge about this time is equivalent to the presence of an external clock), so we have to record all events taking place in selected points of time t_1 and t_2 . From the experimental point of view it means measurements of coincidences between the detectors 1-3 and 2-3 for all possible thicknesses of the birefringent plates A [21, 26, 20]. Averaging the number of counts from detectors 1-3 and 2-3 over the total number of photon pairs from all detectors 1-2-3-4, gives the probability of finding the upper photon in polarization state $|V\rangle$ $p(t_1)$ and $p(t_2)$ at two points of time.

To obtain a more interesting clock, we perform the same conditional probability measurement introducing varying time delays to the clock photon, implemented through quartz plates of variable thickness (dashed box B in Figure 1, block b). [Even though he has no access to abstract coordinate time, he can have access to systems that implement known time delays, that he can calibrate separately.] Now, we obtain a sequence of time-dependent values for the conditional probability: $p(t_1 + \tau_i) = P_{31}^{\tau_i}$ and $p(t_2 + \tau_i) = P_{32}^{\tau_i}$, where $\tau_i = \delta_i/\omega$ is the time delay of the clock photon obtained by inserting the quartz plate B with thickness δ_i in the clock photon path. The experimental results are presented in Fig. 2, where each colour represents a different delay: the yellow points refer to τ_0 ; the red points to τ_1 , etc.

Our results are in very good agreement with the theory (dashed line) [37]. The reduction in visibility of the sinusoidal time dependence of the probability is caused by the decoherence effect due to the use of a low-resolution clock (our clock outputs only two possible values), a well known effect [31, 26, 32].

"Super-observer" mode: The former part of the experiment is addressed to show that one photon "evolves" when using the other photon as a clock. Now we have to show that the global state of two photons is static with respect to any "external" clock. To do so, we need to repeat the whole experiment in "super-observer" mode, and show that for arbitrary thicknesses of the birefringent plates which represent an "external" time the state is invariant. This setup of the experiment is shown in Figure 1(block b). Here the 50/50 beam splitter (BS) on the output of a balanced interferometer is performing a quantum erasure of the polarization measurement performed by the polarizing beam splitter PBS1. In the ideal case the beam splitter should coherently transmit a photon that passes in the upper arm and reflects a photon that passes in the lower arm of the interferometer without affecting its polarization. Without loss of generality, in our setup we use a conventional beam splitter and post-selection on one of the exits. For temporal stability the interferometer was placed into a closed box and narrow interference filters (702 nm, FWHM=1 nm) were inserted before the detectors. A set of quartz plates (A) with different thicknesses are used to introduce an "external" time.

For analyzing an output state of transformed ququarts we used the quantum state tomography procedure. Necessary projective measurements were realized by polarization filters placed in




Figure 2. Experimental results. Probability $p(t)$ that the upper photon is V as a function of the time t recovered from the lower photon. The points with matching colors represent $p(t_1 + \tau_i)$ and $p(t_2 + \tau_i)$: yellow, red, blue, etc., for $i = 0, 1, 2, \dots$, respectively. Here nine different values of τ_i are obtained from a thick quartz plate rotated by nine different angles. The dashed line is the theoretical value.

Table 1. Bell state reconstruction. The table contains values of "external" time and fidelities.

"external" time	Fidelity
0	0.953
$\pi/2\omega$	0.928
$\pi/4\omega$	0.940
$3\pi/2\omega$	0.932

front of detectors. Each filter consists of a sequence of quarter- and half-wave plates and a polarization prism which transmits vertical polarization. Two APDs linked to a coincidence scheme with 1.5 ns time window were used as single photon detectors. In our experiment we used the protocol of Ref. [15, 39]. Registering the coincidence rate for 16 different projections, that were realized by half and quarter plates and a fixed analyzer, it was possible to reconstruct the arbitrary polarization state of ququarts.

The accuracy of quantum tomography is estimated through the fidelity $F = \left(\text{Tr} \sqrt{\rho_{in}^{1/2} \rho_{out} \rho_{in}^{1/2}} \right)^2$ where ρ_{in} is the initial (theoretical) density matrix and ρ_{out} is the reconstructed density matrix after transformation. Our results are summarized in Table I, demonstrating the stationarity of the global state respect to the evolution.

3. Conclusions

In summary, by running our experiment in two different modes ("observer" and "super-observer" mode) we have experimentally shown how the same energy-entangled Hamiltonian eigenstate can be perceived as evolving by the internal observers that test the correlations between a clock subsystem and the rest (also when considering two-time measurements), whereas it is static for the super-observer that tests its global properties. Our experiment is a practical implementation of the Page and Woiters[19] and Gambini et al. [21] mechanisms but, obviously, it cannot discriminate between this and other proposed solutions for the problem of time [4, 5, 6, 2, 7], representing only an "illustration" of this phenomenon. Developments to higher dimensional systems could eventually be envisaged by using atoms [41].

In closing, we note that the time-dependent graphs of Fig. 2 have been obtained without any reference to an external time (or phase) reference, but only from measurements of correlations between the clock photon and the rest representing an implementation of a 'relational' measurement of a physical quantity (time) relative to an internal quantum reference frame

[33, 34], i.e. an example of relational metrology.

Acknowledgments

This research was supported by the John Templeton Foundation (the opinions expressed in this publication are those of the authors and do not necessarily reflect the views of the John Templeton Foundation), E.V.Moreva acknowledges the support from the Dynasty Foundation and Russian Foundation for Basic Research (project 13-02-01170-a).

References

- [1] Aurelius Augustinus Hipponensis Quid est ergo tempus? si nemo ex me quaerat, scio; si quaerenti explicare velim, nescio Confessiones, XI, 14
- [2] Anderson E 2012 *Ann. Phys.* **524** 757 (*Preprint* arXiv:gr-qc/1009.2157v3)
- [3] Sorkin R D 1994 *Int. J. Theor. Phys.* **33** 523; Kiefer C 2009 *Preprint* arXiv:gr-qc/0909.3767; Vedral V 2014 *Preprint* arXiv:quant-ph/1408.6965
- [4] Kuchař K V 1992 Time and interpretations of quantum gravity *Proceedings of the 4th Canadian Conference on General Relativity and Relativistic Astrophysics* (Singapore: World Scientific) 1-92
- [5] Isham C J 1993 Canonical quantum gravity and the problem of time *Integrable Systems, Quantum Groups and Quantum Field Theories* (Dordrecht: Kluwer)
- [6] Ashtekar A 2005 Gravity and the quantum *New J. Phys.* **7** 198
- [7] Merali Z 2013 The origins of space and time *Nature* **500** 516-519
- [8] Nikolic H 2014 *International Journal of Quantum Information* in press, DOI: 10.1142/S0219749915600011; *do.* 2012 *International Journal of Quantum Information* **10** 1250024
- [9] Nicolis S 2014 *Preprint* arXiv:quant-ph/1411.7293
- [10] Briggs J S and Rost J M 2001 *Found. Phys.* **31** 693; Pegg D T 1991 *J. Phys. A* **24** 3031
- [11] Aharonov Y, Popescu S and Tollaksen J 2013 *Preprint* arXiv:quant-ph/1305.1615
- [12] Wootters W K 1984 *Int. J. Theor. Phys.* **23** 701
- [13] Page D N and Wootters W K 1983 *Phys. Rev. D* **27** 2885
- [14] Vedral V and Mott N 1931 *Proc. Camb. Phil. Soc.* **27** 553; Banks T 1985 *Nucl. Phys. B* **249** 332; Brout R 1987 *Found. Phys.* **17** 603; Brout R, Horwitz G and Weil D 1987 *Phys. Lett. B* **192** 318; Brout R 1987 *Z. Phys. B* **68** 339
- [15] James D F V, Kwiat P G, Munro W J and White A G 2001 *Phys. Rev. A* **64** 052312
- [16] Kwiat P, Waks E, White A, Appelbaum I and Eberhard P 1991 *Phys. Rev. A* **60** 773(R)
- [17] DeWitt B S 1967 Quantum theory of gravity. I. the canonical theory *Phys. Rev.* **160** 1113
- [18] Hartle J B and Hawking S W 1983 Wave function of the Universe' *Phys. Rev. D* **28** 2960
- [19] Page D N and Wootters W K 1983 Evolution without evolution: dynamics described by stationary observables *Phys. Rev. D* **27** 2885
- [20] Page D N 1993 Clock time and entropy *Physical Origins of Time Asymmetry* (Cambridge: Cambridge Univ. Press)
- [21] Gambini R, Porto R A, Pullin J and Tortorolo S 2009 Conditional probabilities with Dirac observables and the problem of time in quantum gravity *Phys. Rev. D* **79** 041501(R)
- [22] Peres A 1980 Measurement of time by quantum clocks *Am. J. Phys.* **48** 552
- [23] Rovelli C 1996 Relational Quantum Mechanics *Int. J. Theor. Phys.* **35** 1637
- [24] Rovelli C 1991 *Phys. Rev. D* **43** 442
- [25] Unruh W G and Wald R M 1989 Time and the interpretation of canonical quantum gravity *Phys. Rev. D* **40** 2598
- [26] Gambini R, Pintos L P G and Pullin J 2011 An axiomatic formulation of the Montevideo interpretation of quantum mechanics *Studies His. Phil. Mod. Phys.* **42** 256
- [27] Einstein A 1905 Zur Elektrodynamik bewegter Körper *Ann. Phys.* **17** 891; translated in *do.*1923 *The Principle of Relativity* (London: Methuen & Co.)
- [28] Scully M O, Englert B G and Walther H 1991 Quantum optical tests of complementarity *Nature* **351** 111
- [29] Dürr S, Nonn T and Rempe G 1998 Origin of quantum-mechanical complementarity probed by a 'which-way' experiment in an atom interferometer *Nature* **395** 33
- [30] Preskill J *Lecture Notes for Physics* 229 (<http://www.theory.caltech.edu/people/preskill/ph229>, p 68)
- [31] Gambini R, Pintos L P G and Pullin J 2010 Undecidability and the problem of outcomes in quantum measurements it *Found. Phys.* **40** 93
- [32] Egusquiza I L, Garay L J and Raya J M 1999 Quantum evolution according to real clocks *Phys. Rev. A* **59** 3236

- [33] Bartlett S D, Rudolph T and Spekkens W 2007 Reference frames, superselection rules, and quantum information *Rev. Mod. Phys.* **79** 555
- [34] Wiseman H M 2004 Defending continuous variable teleportation: why a laser is a clock, not a quantum channel *J. Opt. B: Quantum Semiclass. Opt.* **6** S849
- [35] Tiefenbacher F *et al* 2007 Entanglement-based quantum communication over 144 km *Nature Physics* **3** 481; Boschi D *et al* 1998 Experimental Realization of Teleporting an Unknown Pure Quantum State via Dual Classical and Einstein-Podolsky-Rosen Channels *Phys. Rev. Lett.* **80** 1121; Kwiat P G *et al* 2001 Experimental entanglement distillation and hidden non-locality *Nature* **409** 1014; Yamamoto T *et al* 2003 Experimental extraction of an entangled photon pair from two identically decohered pairs *Nature* **421** 343; Pan J W *et al* 2001 Entanglement purification for quantum communication *Nature* **410** 1067; Brida G *et al* 2012 *Phys. Rev. A* **68**, 033803; Brida G *et al* 2006 *Phys. Rev. Lett.* **96** 143601
- [36] Genovese M 2005 *Phys. Rep.* **413** 319
- [37] Moreva E V, Brida G, Gramegna M, Giovannetti V, Maccone L and Genovese M 2014 *Phys. Rev. A* **89** 052122
- [38] Bogdanov Yu I, Brida G, Genovese M, Shurupov A P, Moreva E V and Kulik S P 2010 *Phys. Rev. Lett.* **105** 010404
- [39] Bogdanov Yu I, Moreva E V, Maslennikov G A, Galeev R F, Straupe S S and Kulik S P 2006 *Phys. Rev. A* **73** 063810
- [40] Brida G, Degiovanni I P, Florio A, Genovese M, Giorda P, Meda A, Paris M and Shurupov A 2010 *Phys. Rev. Lett.* **104** 100501
- [41] Massar S, Spindel P, Varn A F and Wunderlichet C 2014 *Preprint* arXiv:quant-ph 1410.6683