



ISTITUTO NAZIONALE DI RICERCA METROLOGICA
Repository Istituzionale

Anomalous weak values via a single photon detection

Original

Anomalous weak values via a single photon detection / Rebufello, E; Piacentini, F; Avella, A; de Souza, Ma; Gramegna, M; Dziewior, J; Cohen, E; Vaidman, L; Degiovanni, Ip; Genovese, M. - In: LIGHT, SCIENCE & APPLICATIONS. - ISSN 2047-7538. - 10:1(2021), p. 106. [10.1038/s41377-021-00539-0]

Availability:

This version is available at: 11696/72890 since: 2022-02-11T10:33:06Z

Publisher:

SPRINGERNATURE

Published

DOI:10.1038/s41377-021-00539-0

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

LETTER

Open Access

Anomalous weak values via a single photon detection

Enrico Rebufello¹, Fabrizio Piacentini¹, Alessio Avella¹, Muriel A. de Souza², Marco Gramegna¹, Jan Dziewior^{3,4}, Eliahu Cohen⁵, Lev Vaidman⁶, Ivo Pietro Degiovanni¹ and Marco Genovese¹

Abstract

Is it possible that a measurement of a spin component of a spin-1/2 particle yields the value 100? In 1988 Aharonov, Albert and Vaidman argued that upon pre- and postselection of particular spin states, weakening the coupling of a standard measurement procedure ensures this paradoxical result¹. This theoretical prediction, called *weak value*, was realised in numerous experiments^{2–9}, but its meaning remains very controversial^{10–19}, since its “anomalous” nature, i.e., the possibility to exceed the eigenvalue spectrum, as well as its “quantumness” are debated^{20–22}. We address these questions by presenting the first experiment measuring anomalous weak values with just a single click, without the need for statistical averaging. The measurement uncertainty is significantly smaller than the gap between the measured weak value and the nearest eigenvalue. Beyond clarifying the meaning of weak values, demonstrating their non-statistical, single-particle nature, this result represents a breakthrough in understanding the foundations of quantum measurement, showing unprecedented measurement capability for further applications of weak values to quantum photonics.

Weak values (WVs), as introduced in ref. ¹, represent one of the most interesting and intriguing quantum measurement paradigms. In that influential work, the outcome of the measurement of the spin component was 100, while in the chosen units its maximal eigenvalue was just 1. However, weakening the coupling in the measurement procedure made the uncertainty in an individual measurement much larger than 1 (and even than 100), thus this “anomalous” value was observed only after averaging over a very large number of readings of the pointer variable. While averaging is a standard practice in many measurement protocols, postselection is not, hence the legitimacy of the statistical analysis was questioned^{20–22}. Understanding this matter is fundamental not only for clarifying the meaning of WVs, but also in view of significant applications in quantum metrology^{3–5,23,24}. In this work, we present a *robust weak measurement*, an experiment in which a single reading of the

measuring device, coupled to the system only once, provides a WV and, in particular, an anomalous one. Postselection still plays a crucial role, but the anomalous outcome no longer arises from a statistical analysis. We measured an observable with eigenvalues in the range $[-7, 7]$. The WV of the observable of the pre- and postselected system on which a single-click measurement was performed was 18.7, and our single click yielded 21.4 ± 4.5 , see Fig. 1.

This is a surprising result, since the expectation value of the observable in the preselected state was only 2.2. It would not be surprising that postselection on the eigenstate corresponding to a maximal eigenvalue (i.e., 7) slightly increased the measured value, but only up to 7, not beyond (in fact, in our experiment the expectation value corresponding to the postselected state was also just 2.2).

The main theoretical basis for our experiment is the work in ref. ²⁵, which preceded the introduction of WVs¹, extended here by constructing a new method allowing a feasible experimental implementation. The easiest way to explain this work is via weak measurements performed on a system consisting of n particles with a single measuring device, see Sec. VII of ref. ²⁶. Within the standard weak

Correspondence: Fabrizio Piacentini (f.piacentini@inrim.it)

¹INRIM, Strada delle Cacce 91, I-10135 Torino, Italy


²National Institute of Metrology, Quality and Technology—INMETRO, Av. Nossa Senhora das Graças, 50, Duque de Caxias, RJ 25250-020, Brazil

Full list of author information is available at the end of the article

© The Author(s) 2021



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.



measurement procedure, in which each particle has its own measuring device, the uncertainty of measuring the sum of variables $A^\Sigma \equiv \sum_{k=1}^n A_k$, where A_k corresponds to particle k , scales like \sqrt{n} because of the contributions of the n measuring devices. In our procedure for measuring A^Σ , we couple a single measuring device to all particles and thus, importantly, the final uncertainty is the one of the single measuring device involved, avoiding the \sqrt{n} increase. This allows extracting an anomalous WV $(A^\Sigma)_w$ by coupling the measuring device only once to each part of the system, and observing only a single click of the measuring device.

In our experimental demonstration, the measured variable is the sum of polarisation variables of n photons:

$$\sigma_3^\Sigma \equiv \sum_{k=1}^n \sigma_3^{(k)} \tag{1}$$

where $\sigma_3 \equiv |H\rangle\langle H| - |V\rangle\langle V|$, and $H(V)$ is the horizontal (vertical) polarisation. All photons are preselected in the state

$$|\psi_\alpha\rangle = \cos \alpha |H\rangle + \sin \alpha |V\rangle \tag{2}$$

and postselected in the state

$$|\psi_\beta\rangle = \cos \beta |H\rangle + \sin \beta |V\rangle \tag{3}$$

When states (2) and (3) are almost orthogonal, we obtain an anomalously large weak value

$$(\sigma_3^\Sigma)_w = \frac{\langle \psi_\beta | \sigma_3^\Sigma | \psi_\alpha \rangle}{\langle \psi_\beta | \psi_\alpha \rangle} \tag{4}$$

All photons are coupled to the same measuring device, so the interaction Hamiltonian is

$$\mathcal{H} = g(t) \sum_{k=1}^n \sigma_3^{(k)} \otimes p_x \tag{5}$$

where a well localized $g(t)$ defines the time of the measurement, $\int g(t) dt = 1$, and p_x is the conjugate momentum of the pointer variable x with the initial quantum state modelled by the Gaussian

$$\chi(x) = \frac{1}{\sqrt{\Delta\sqrt{2\pi}}} e^{-x^2/4\Delta^2} \tag{6}$$

The strength of the measurement interaction in Eq. (5) is characterized by $\frac{1}{\Delta}$, which has to be chosen carefully: for a single-click measurement the uncertainty cannot be too large, but also the coupling cannot be too strong, as it changes the WV^{19,27}. Moreover, although the system has a well-defined WV of the measured observable at every moment between pre- and postselection, the measuring device, in general, does not indicate this WV, because of entanglement between system and measuring device, see Fig. 5 in ref. ²². Somewhat surprisingly, in the case of coupling to a Gaussian pointer, the expectation value of the pointer after the measurement exactly equals this WV. Since, in the present scenario, the WV is constant in time during the measurement interaction, we can extract it at the moment just before the postselection, by calculating the quantum state after the interaction and knowing the postselection state.

To make the explanation more transparent, let us first consider a case for $n = 1$. The state of the photon and the measuring device before the postselection is

$$\cos \alpha |H\rangle |\chi_+\rangle + \sin \alpha |V\rangle |\chi_-\rangle \tag{7}$$

where $|\chi_+\rangle$ and $|\chi_-\rangle$ denote Gaussians shifted by 1 and -1 , respectively. Thus, the photon has a mixed polarisation state described by the density matrix ρ_α that can be expressed in the $\{|H\rangle, |V\rangle\}$ basis as

$$\rho_\alpha = \begin{pmatrix} \cos^2 \alpha & e^{-\frac{1}{2\Delta^2}} \sin \alpha \cos \alpha \\ e^{-\frac{1}{2\Delta^2}} \sin \alpha \cos \alpha & \sin^2 \alpha \end{pmatrix} \tag{8}$$

Postselecting on $|\psi_\beta\rangle$, the WV is given by (see Eq. (32) in ref. ²²)

$$(\sigma_3)_w = \frac{\text{tr}(|\psi_\beta\rangle\langle\psi_\beta| \sigma_3 \rho_\alpha)}{\text{tr}(|\psi_\beta\rangle\langle\psi_\beta| \rho_\alpha)} = \frac{\mu^2 - \nu^2}{\mu^2 + \nu^2 + 2\mu\nu e^{-\frac{1}{2\Delta^2}}} \tag{9}$$

where $\mu = \cos \alpha \cos \beta$ and $\nu = \sin \alpha \sin \beta$.

It is straightforward to generalise the calculation for $n > 1$. We define the joint pre- and postselected states of the n photons as, respectively, $|\Psi_\alpha\rangle = \bigotimes_{k=1}^n |\psi_\alpha^{(k)}\rangle$ and $|\Psi_\beta\rangle = \bigotimes_{k=1}^n |\psi_\beta^{(k)}\rangle$. The WV is


$$\begin{aligned} (\sigma_3^\Sigma)_w &= \frac{\langle \Psi_\beta | \sigma_3^\Sigma \text{tr}(U^\Sigma |\Psi_\alpha\rangle \langle \Psi_\alpha| \otimes |x\rangle \langle x| (U^\Sigma)^\dagger) | \Psi_\beta \rangle}{\langle \Psi_\beta | \text{tr}(U^\Sigma |\Psi_\alpha\rangle \langle \Psi_\alpha| \otimes |x\rangle \langle x| (U^\Sigma)^\dagger) | \Psi_\beta \rangle} \\ &= \frac{\sum_{k,l=0}^n \binom{n}{k} \binom{n}{l} \mu^{k+l} \nu^{2n-k-l} (2k-n) \gamma_{kl}}{\sum_{k,l=0}^n \binom{n}{k} \binom{n}{l} \mu^{k+l} \nu^{2n-k-l} \gamma_{kl}} \end{aligned} \quad (10)$$

where $U^\Sigma = e^{-i \sum_{k=1}^n \sigma_3^{(k)} \otimes p_x}$, $\gamma_{kl} = e^{-\frac{(k-l)^2}{2\Delta^2}}$, and the trace is taken over the pointer system only. Our Gaussian pointer, after the measurement, shows $\langle x \rangle = (\sigma_3^\Sigma)_w$ with uncertainty $\Delta x = \sqrt{\langle x^2 \rangle - \langle x \rangle^2}$, where

$$\langle x^2 \rangle = \frac{\sum_{k,l=0}^n \binom{n}{k} \binom{n}{l} \mu^{k+l} \nu^{2n-k-l} ((n-k-l)^2 + \Delta^2) \gamma_{kl}}{\sum_{k,l=0}^n \binom{n}{k} \binom{n}{l} \mu^{k+l} \nu^{2n-k-l} \gamma_{kl}} \quad (11)$$

The experiment directly testing this prediction, see Fig. 2a, is very difficult.

The following observation leads to a more feasible test of our predictions. In the Hamiltonian in Eq. (5) the variables $\sigma_3^{(k)}$ are constants of motion, therefore it does not matter when the interaction with every photon takes place. In particular, we can perform the couplings one



after another. In this case, however, we can use just one photon instead of n , pre- and postselecting it n times according to the states in Eqs. (2) and (3), see Fig. 2b. The observable, i.e., the sum of polarisation variables of n photons, is replaced by the sum of polarisation variables of the same photon at n different times, which is much easier to measure, since the spatial degree of freedom of this single photon can play the role of the measuring device.

We consider the case $n = 7$, which is large enough to see the predicted effect. In Fig. 3 we show the WV $(\sigma_3^\Sigma)_w$ and the pointer uncertainty Δx (Eqs. (10) and (11), respectively) for $\alpha = 0.62$ and $\Delta = 5.84$, as a function of β . By choosing $\beta = 2.53$, one obtains a theoretical final state of the pointer showing 18.7 ± 4.5 , granting an optimal reading of the anomalous WV in terms of relative uncertainty. Note that the final width of the pointer is smaller than the initial width, due to a subtle narrowing effect²⁸.

The experimental setup is shown in Fig. 4. It is composed of a set of $n = 7$ blocks in which a birefringent crystal pair realises the weak interaction, preceded by a half-wave plate and followed by a polarising plate. While the polarising plate performs the postselection, the half-wave plate rotates the polarisation of the photon outgoing the previous block to set the preselection state. The EM-CCD placed at the end of the $n = 7$ blocks detects the arrival position of the photon. A detailed description of the setup is provided in the Methods section.

The single-click experiment presented in Fig. 1 yielded 21.4, in agreement with the predicted value within the theoretical uncertainty.

We tested our theoretical analysis a posteriori by performing a multi-click experimental run with the same parameters α , β and Δ , to record the distribution shown in Fig. 5. The mean value of this multi-click distribution, 18.59, is very close to the theoretical WV $(\sigma_3^\Sigma)_w = 18.7$, given by Eq. (10). The statistical uncertainty is only 0.09, thus the remaining discrepancy is due to imprecision in the parameters α , β , Δ fixing the theoretical value. Some deviations are caused by imperfections in the optical elements, e.g., the birefringent crystals (details in the Supplementary Material). The width of the multi-click distribution in Fig. 5 turns out to be 4.5, in full agreement with the theoretical predictions.

As in all weak measurements in which the pointer variable of the polarisation measurement of a photon is its transverse degree of freedom, the probability distribution in Fig. 5 is identical to the intensity pattern of the electromagnetic wave on the screen which can be calculated using Maxwell equations (see refs. 4,29). This is a sufficiently narrow distribution to allow a robust measurement of the WV via the detection of a single quantum particle. In the framework of classical physics, this bizarre result is hidden in the peculiar interference phenomena of




Fig. 3 Predicted weak value and pointer uncertainty. Predicted values for the WV $(\sigma_3^\Sigma)_w$ (solid blue line) and the final pointer uncertainty Δx (solid red line) for $n = 7$, $\alpha = 0.62$ and $\Delta = 5.84$, as a function of β . The initial photon distribution width Δ is included as a dashed brown line. The two black dots on the curves denote the $(\sigma_3^\Sigma)_w = 18.7$ and $\Delta x = 4.5$ obtained for $\beta = 2.53$, the parameters chosen for our experimental demonstration




Fig. 4 Experimental setup. Our photon source exploits type-I spontaneous parametric down conversion. Generated signal photons at 702 nm are spectrally-filtered, injected in a single-mode fibre and then collimated in a Gaussian mode to be used in the experiment, while idler photons at 920 nm are detected by a Single-Photon Avalanche Detector in order to monitor the stability of the source. The robust weak measurement is obtained by means of the $n = 7$ identical blocks put after the initial PBS. A spatially-resolving detector (EM-CCD camera operating in the photon counting regime) is used to determine the final position of the detected photons

superoscillations^{30,31}. Thus, the quantum formalism of robust weak values, apart from describing a genuinely quantum effect for the case of measurement with an external pointer, naturally incorporates this classical interference effect.

To increase confidence in our results, we repeated the experiment for a few sets of different parameters leading to less anomalous WVs (and even a non-anomalous one). All results are presented in Table 1 (see the Supplementary Material for additional information). The results are influenced by uncertainties originating from the calibration procedure. However, when the purpose is not to find a precise numerical value of the polarisation WV, but to

test the single-click measurement method versus an ensemble measurement, the calibration uncertainty is irrelevant. The experimental data shown in Table 1 fit the theoretical predictions well, proving the possibility of measuring an anomalous WV with a single detection.

Theoretical calculations, see Fig. 3, show that the final uncertainty of the pointer is close to the initial beam width Δ . Thus, faced with a new task of estimating $(\sigma_3^\Sigma)_w$ in which somebody else fixes the pre- and postselected states of the system, a single click on our detector is capable of providing the WV with an uncertainty of the same order of magnitude as the width of the initial beam even for anomalous WVs.

In summary, our results offer a deeper understanding of the meaning of WVs, providing a significant contribution to the development of quantum measurement in the weak coupling regime. Estimating anomalous WVs with a single-click breaks the common belief that the WV is a statistical concept of conditional expectation value¹³. Our findings, going well beyond refs. 22,32, stress the non-statistical, single-particle nature of WVs, demonstrating how a single-click measurement can provide a WV estimate even for anomalous WVs. Furthermore, this experiment suggests a viable possibility for amplification methods effectively reducing the uncertainty contribution associated with the measurement of the pointer. This paves the way for future practical applications of the robust weak measurement paradigm.

Methods

In our experimental setup, shown in Fig. 4 (further details in the Supplementary Material), photons in a multi-thermal distribution with a mean photon number per pulse $\ll 1$ are produced by type-I spontaneous parametric down-conversion (SPDC). This guarantees a short coherence time (~ 150 fs), avoiding unwanted self-interference effects due to internal reflections. The SPDC occurs in a $10 \times 10 \times 5$ mm LiIO₃ crystal, pumped by the second harmonic generation (398 nm) of a 76 MHz mode-locked laser at 796 nm. The signal photons are spectrally filtered and coupled to a single-mode fibre. At the end of the fibre, the photons are collimated in a Gaussian mode and sent to the free-space path where the robust weak measurement experiment occurs. After passing through an initial polarising beam splitter (PBS), used to suppress any residual circular polarisation component, the signal photons go through $n = 7$ identical blocks, each implementing three steps: preselection, weak coupling and postselection. Each photon enters every block in a linear polarisation state, due to either the first PBS or the postselection of the previous block. Every block begins with a quartz half-wave plate, implementing a unitary rotation R aligning the photon

polarisation axis to the direction corresponding to the initial (preselected) state $|\psi_\alpha\rangle$. Then, a birefringent unit composed of a pair of birefringent crystals is responsible for the weak interaction. The first calcite crystal, 2 mm long, has the extraordinary (e) optical axis lying in the $x - z$ plane, forming an angle of $\pi/4$ with respect to the z direction. This generates a spatial walk-off (of ~ 0.2 mm) along the x direction for the horizontally-polarised photons, reducing the overlap between the horizontal and vertical polarisation components. The second crystal of each unit is a 1.1 mm long calcite crystal with the optical e -axis lying along the y direction. It generates no spatial walk-off, and its role is to compensate the temporal walk-off induced by the first crystal. The last component of each block is a polarising plate, postselecting the photons in the state $|\psi_\beta\rangle$. After the $n = 7$ blocks, the photons are detected




Table 1 Results for the various parameters of the measurement setup

#	1	2	3	4	5	6	7	8	9	10
	α	β	Δ	$(\sigma_3^\Sigma)_w$	$\Delta[(\sigma_3^\Sigma)_w]$	$(\sigma_3^\Sigma)_{w}^{\text{exp}}$	$\Delta_{\text{stat}}[(\sigma_3^\Sigma)_{w}^{\text{exp}}]$	$(\sigma_3^\Sigma)_w^{\text{1click}}$	$\Delta(\sigma_3^\Sigma)_w^{\text{1click}}$	Δx
(a)	0.62	2.53	5.84	18.7	0.9	18.59	0.09	21.4	4.5	4.5
(b)	0.62	2.53	3.18	9.8	0.8	10.51	0.02	10.9	3.8	5.0
(c)	0.52	2.62	2.96	11.4	0.5	11.07	0.08	14.1	3.4	2.8
(d)	0.52	0.88	3.09	1.3	0.4	0.97	0.05	-2.6	4.6	3.6

Columns 1-4 describe the preparation parameters and the corresponding weak value $(\sigma_3^\Sigma)_w$. Column 5 shows the systematic uncertainty in the experimental implementation of $(\sigma_3^\Sigma)_w$ due to the uncertainties in the preparation parameters α , β and Δ (which incorporate inhomogeneities in birefringent crystals and other experimental imperfections). Column 6 shows the experimental mean values obtained by repeating the single-photon experiments (EM-CCD dark counts subtracted). The statistical uncertainty in these experiments is shown in column 7. Column 8 presents the experimental weak values $(\sigma_3^\Sigma)_w^{\text{1click}}$ extracted from a single detection event. The uncertainty in column 9 represents the quantum uncertainty of the pointer variable experimentally obtained from repeated measurements with the same parameters as the single-click experiment (see histogram in Fig. 5). Column 10 contains the predicted final uncertainty of the pointer, Δx , calculated from the parameters α , β , and Δ

by a spatially resolving detector, i.e., an Electron Multiplying CCD (EM-CCD) device able to work in the linear analogue regime as well as in the photon counting regime (details in ref. ³³). To calibrate our system, we identify the extremal positions of the pointer on our EM-CCD, i.e. the position of the $|V\rangle$ polarisation state, corresponding to the eigenvalue $\sigma_3^\Sigma = -7$ and then the position of the state $|H\rangle$, corresponding to $\sigma_3^\Sigma = 7$. This allows us to define the zero point as the centre between the readings for $\sigma_3^\Sigma = \pm 7$, and also scale the pointer variable accordingly.

Acknowledgements

We acknowledge the European Union's Horizon 2020 and the EMPIR Participating States in the context of the projects 17FUN01 "BeCOMe" and 17FUN06 "SIQUST", the European Union's Horizon 2020 FET-OPEN project grant no. 828946 "PATHOS", the National Science Foundation—U.S.-Israel Binational Science Foundation Grant No. 735/18. EC was supported by Grant No. FQXi-RFP-CPW-2006 from the Foundational Questions Institute and Fetzer Franklin Fund, a donor advised fund of Silicon Valley Community Foundation, the Israel Innovation Authority under Projects No. 70002 and No. 73795, the Quantum Science and Technology Program of the Israeli Council of Higher Education, and the Pazy Foundation. JD acknowledges support from the PhD program IMPRS-QST.

Author details

¹INRIM, Strada delle Cacce 91, I-10135 Torino, Italy. ²National Institute of Metrology, Quality and Technology—INMETRO, Av. Nossa Senhora das Graças, 50, Duque de Caxias, RJ 25250-020, Brazil. ³Max-Planck-Institut für Quantenoptik, Hans-Kopfermann-Straße 1, 85748 Garching, Germany. ⁴Department für Physik, Ludwig-Maximilians-Universität, 80797 München, Germany. ⁵Faculty of Engineering and the Institute of Nanotechnology and Advanced Materials, Bar Ilan University, Ramat Gan 5290002, Israel. ⁶Raymond and Beverly Sackler School of Physics and Astronomy, Tel-Aviv University, Tel-Aviv 6997801, Israel

Author contributions

I.P.D., M.Gram, M.Gen (responsible of the laboratories) and L.V. (responsible for the theoretical framework) planned the experiment, with the support of F.P., A.A. and E.C. The experimental realisation was achieved (supervised by IPD, M.Gram and M.Gen) by ER (leading role), F.P., A.A. and M.A.de.S. The theoretical aspects were addressed by J.D., E.C. and L.V.. The manuscript was prepared with inputs by all the authors. They also had a fruitful systematic discussion on the progress of the work.

Conflict of interest

The authors declare no competing interests.

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41377-021-00539-0>.

Received: 9 October 2020 Revised: 7 April 2021 Accepted: 19 April 2021
Published online: 25 May 2021

References

- Aharonov, Y., Albert, D. Z. & Vaidman, L. How the result of a measurement of a component of the spin of a spin-1/2 particle can turn out to be 100. *Phys. Rev. Lett.* **60**, 1351–1354 (1988).
- Ritchie, N. W. M., Story, J. G. & Hulet, R. G. Realization of a measurement of a "weak value". *Phys. Rev. Lett.* **66**, 1107–1110 (1991).
- Hosten, O. & Kwiat, P. Observation of the spin hall effect of light via weak measurements. *Science* **319**, 787–790 (2008).
- Dixon, P. B. et al. Ultrasensitive beam deflection measurement via interferometric weak value amplification. *Phys. Rev. Lett.* **102**, 173601 (2009).
- Xu, X. Y. et al. Phase estimation with weak measurement using a white light source. *Phys. Rev. Lett.* **111**, 033604 (2013).
- Piacentini, F. et al. Experiment investigating the connection between weak values and contextuality. *Phys. Rev. Lett.* **116**, 180401 (2016).
- Lundeen, J. S. et al. Direct measurement of the quantum wavefunction. *Nature* **474**, 188–191 (2011).
- Goggin, M. E. et al. Violation of the Leggett-Garg inequality with weak measurements of photons. *Proc. Natl. Acad. Sci. USA* **108**, 1256–1261 (2011).
- Magaña Loaiza, O. et al. Amplification of angular rotations using weak measurements. *Phys. Rev. Lett.* **112**, 200401 (2014).
- Leggett, A. J. Comment on "How the result of a measurement of a component of the spin of a spin-1/2 particle can turn out to be 100". *Phys. Rev. Lett.* **62**, 2325 (1989).
- Peres, A. Quantum measurements with postselection. *Phys. Rev. Lett.* **62**, 2326 (1989).
- Aharonov, Y. & Vaidman, L. Aharonov and Vaidman reply. *Phys. Rev. Lett.* **62**, 2327 (1989).
- Dressel, J. et al. Colloquium: understanding quantum weak values: basics and applications. *Rev. Modern Phys.* **86**, 307 (2014).
- Ferrie, C. & Combes, J. Weak value amplification is suboptimal for estimation and detection. *Phys. Rev. Lett.* **112**, 040406 (2014).
- Jordan, A. N., Martínez-Rincón, J. & Howell, J. Technical advantages for weak-value amplification: when less is more. *Phys. Rev. X* **4**, 011031 (2014).
- Pang, S. S., Dressel, J. & Brun, T. A. Entanglement-assisted weak value amplification. *Phys. Rev. Lett.* **113**, 030401 (2014).
- Pusey, M. F. Anomalous weak values are proofs of contextuality. *Phys. Rev. Lett.* **113**, 200401 (2014).
- Vaidman, L. Weak value controversy. *Philos. Trans. R Soc. A, Math. Phys. Eng. Sci.* **375**, 21060395 (2017).
- Piacentini, F. et al. Investigating the effects of the interaction intensity in a weak measurement. *Sci. Rep.* **8**, 6959 (2018).
- Ferrie, C. & Combes, J. How the result of a single coin toss can turn out to be 100 heads. *Phys. Rev. Lett.* **113**, 120404 (2014).
- Mundarain, D. F. & Orszag, M. Quantumness of the anomalous weak measurement value. *Phys. Rev. A* **93**, 032106 (2016).
- Vaidman, L. et al. Weak value beyond conditional expectation value of the pointer readings. *Phys. Rev. A* **96**, 032114 (2017).
- Hallaji, M. et al. Weak-value amplification of the nonlinear effect of a single photon. *Nat. Phys.* **13**, 540–544 (2017).
- Lyons, K. et al. Power-recycled weak-value-based metrology. *Phys. Rev. Lett.* **114**, 170801 (2015).
- Aharonov, Y. et al. Surprising quantum effects. *Phys. Lett. A* **124**, 199–203 (1987).
- Aharonov, Y. & Vaidman, L. Properties of a quantum system during the time interval between two measurements. *Phys. Rev. A* **41**, 11 (1990).
- Dziwior, J. et al. Universality of local weak interactions and its application for interferometric alignment. *Proc. Natl. Acad. Sci. USA* **116**, 2881–2890 (2019).
- Aharonov, Y. & Botero, A. Quantum averages of weak values. *Phys. Rev. A* **72**, 052111 (2005).
- Howell, J. C. et al. Interferometric weak value deflections: quantum and classical treatments. *Phys. Rev. A* **81**, 033813 (2010).
- Berry, M. V. & Popescu, S. Evolution of quantum superoscillations and optical superresolution without evanescent waves. *J. Phys. A: Math. Gen.* **39**, 6965–6977 (2006).
- Aharonov, Y. et al. Some mathematical properties of superoscillations. *J. Phys. A: Math. Theor.* **44**, 365304 (2011).
- Piacentini, F. et al. Determining the quantum expectation value by measuring a single photon. *Nat. Phys.* **13**, 1191–1194 (2017).
- Avella, A. et al. Absolute calibration of an EMCCD camera by quantum correlation, linking photon counting to the analog regime. *Opt. Lett.* **41**, 1841–1844 (2016).