

ISTITUTO NAZIONALE DI RICERCA METROLOGICA
Repository Istituzionale

Counting atoms

This is the author's accepted version of the contribution published as:

Original

Counting atoms / Massa, Enrico; Mana, Giovanni. - In: NATURE PHYSICS. - ISSN 1745-2473. - 12:5(2016), pp. 522-522. [10.1038/nphys3754]

Availability:

This version is available at: 11696/53079 since: 2020-05-21T11:15:13Z

Publisher:

Nature Publishing Group

Published

DOI:10.1038/nphys3754

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Physical sciences / Physics [URI /639/766]

Physical sciences / Chemistry / Analytical chemistry [URI /639/638/11]

Physical sciences / Physics / Techniques and instrumentation / Characterization and analytical techniques [URI /639/766/930/12]

Social science / Science, technology and society [URI /4014/4045]

Scientific community and society [URI /706]

Counting atoms

Enrico Massa and Giovanni Mana expound on the substance of the Avogadro constant.

The need for a physical constant quantifying a reference amount of substance is a manifestation of the atomic nature of matter. At the basis of the concept of a 'quantity constant' lies the law of constant proportions: a pure compound always contains the same elements — the conserved quantities in chemical reactions — in the same mass ratio.

This law played a fundamental part in the metrological development of chemistry: the amount of a given element or compound can be traced back to the amount of substance in 12 g of ^{12}C — a mole of ^{12}C — by chains of stoichiometric proportions. For example, since 12 g of ^{12}C reacts with (approximately, ignoring less abundant isotopes) 4 g of H_2 and $\text{C} + 2\text{H}_2 \rightarrow \text{CH}_4$, one mole of molecular hydrogen corresponds to 2 g. Next, since 2 g of H_2 reacts with 16 g of O_2 and $\text{O}_2 + 2\text{H}_2 \rightarrow 2\text{H}_2\text{O}$, the molar mass of molecular oxygen is 32 g.

In the 19th century, investigations of a number of chemists brought to light clues that 'amount of substance' is discrete: it always comes in integer quanta, basic amounts that take part in chemical reactions. These 'basic amounts' were found to be constant — independent of substance, time, place or temperature, This strengthened the atomic hypothesis and linked the vindication of the existence of atoms to determining the number of quanta in a mole, the Avogadro constant N_{A} , introduced in 1909 by Jean Perrin and named after Amedeo Avogadro, who pioneered the idea of molecules.

The original reference substance was chosen to be hydrogen — in modern parlance, the mole was set to the number of atoms in 1 g of ^1H . Later, oxygen was used, because it forms more compounds. Eventually, in 1960, ^{12}C was chosen.

Although the concept of the atom goes back to the Greek philosopher Democritus, there was no experimental proof of atomism until the end of the 19th century. One remarkable attempt to count the atoms in a finite amount of substance is that of Johann Magnenus, who reported in 1646 that the scent of burned incense fills a church of more than seven hundred million times its original size and that this observation can be used to arrive at a primitive estimate of an atom's size [1]. From the ratio of the volumes of the interior of the church and the minimal amount of evenly dispersed incense that one can sense, which he estimated to be to one thousandth of a chickpea and guessed to contain at least a million atoms, Magnenus arrived at 7.776×10^{17} as a lower bound to the number of atoms in an incense chickpea. What he did — although not very reliably — is the same we do these days

to count atoms in silicon crystals. Both approaches ingeniously amplify the atomic scale so that macro-scale detection is possible: Magnenus used diffusion and smelling, we use crystallization and X-ray diffraction.

From 1865 to 1965 more than eighty determinations of N_A were reported, obtained by means of more than twenty different methods [2]; notable cases include those by Loschmidt (kinetic theory of gases), Perrin (Brownian motion) and Einstein (diffusion of particles in a fluid).

Up to 1965, when Bonse and Hart demonstrated the first X-ray interferometer [3], the value of N_A was used for determining the lattice spacings in crystals from their volumes and masses. Now, X-ray interferometry enables counting how many silicon atoms are aligned in a crystal the length of which is measured by optical interferometry and to relate atomic dimensions to the wavelength of visible light [4]. Using this method, the number of atoms in a 1 kg sphere of high-purity ^{28}Si (an artist's impression is pictured, featuring Avogadro and his law) was counted and N_A determined by dividing the sphere's volume by ^{28}Si 's atomic volume measured via X-ray interferometry. The currently most accurately measured value of N_A , $6.02214076 \times 10^{23} \text{ mol}^{-1}$ with a relative standard uncertainty of 20 parts in 10^9 , has been obtained in this way by the International Avogadro Coordination project [5].

High-precision measurements of the Avogadro constant are important for being able to quantitatively connect the macro- and microscales — obviously, knowing N_A , one can switch between molar mass and particle mass. But in addition, metrologists are planning to define the kilogram and the mole via the Planck and Avogadro constants. Before that can happen, an accurate determination of N_A needs to be realized. By fixing N_A , the atomic model of matter becomes embedded in the International System of Units (SI) and the mole redefined in terms of the number of entities it represents. Furthermore, since the ratio between the ^{28}Si mass and the Planck constant can be accurately measured via atom interferometry, and since we are now able to 'count atoms', macroscopic prototypes with an exactly known amount of substance and mass can be realized — neglecting Einstein's mass–energy equivalence principle and atoms therefore having unique and well-defined masses...

Enrico Massa and Giovanni Mana are at the Istituto Nazionale di Ricerca Metrologica, Strada delle Cacce 91, 10135 Torino, Italy.
E-mail: e.massa@inrim.it, g.mana@inrim.it

References

- [1] Johann Chrysostom Magnenus 1646 *Democritus reviviscens* 207
- [2] Mana G and Zosi G 1995 *Rivista del Nuovo Cimento* 18 1
- [3] Bone U and Hart M 1968 *Appl. Phys. Lett.* **6** 155
- [4] Massa E *et al.* 2015 *J. Phys. Chem. Ref. Data* **44** 031208
- [5] Azuma Y *et al.* 2015 *Metrologia* **52** 360-375



Courtesy of Angelo Bergamin
bergamin.a@tiscali.it